

Retrospective Assessment of Environmental Pollution by PBDEs and HBCDs Using Sediment Core from Manila Bay, the Philippines

Tomohiko ISOBE^{1,2}, Atsuko AMANO³, Kwang-Hyeon CHANG¹,
Rommel H. MANEJA⁴, Peter B. ZAMORA⁴,
Maria Lourdes SAN DIEGO-MCGLONE⁴, Fernando P. SIRINGAN⁴,
Maricar PRUDENTE⁵, Todd W. MILLER¹ and Shinsuke TANABE¹

¹*Center for Marine Environmental Studies (CMES), Ehime University,
2-5 Bunkyo-cho, Matsuyama 790-8577, Japan*

²*Senior Research Fellow Center, Ehime University,
2-5 Bunkyo-cho, Matsuyama 790-8577, Japan*

³*Institute of Geology and Geoinformation,
AIST, 1-1-1 Higashi, Tsukuba 305-8561, Japan*

⁴*Marine Science Institute, University of the Philippines,
Diliman, Quezon City 1101, Philippines*

⁵*Science Education Department, De La Salle University, Manila, Philippines*

(Received 21 March 2010; accepted 6 June 2010)

Abstract—Brominated flame retardants, including polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecanes (HBCDs), as well as PCBs were determined in surface and core sediments from Manila Bay, the Philippines. Organohalogen compounds analyzed in the present study were detected in most of the surface sediment samples, indicating ubiquitous pollution by these contaminants. Concentrations of BDE-209 were slightly higher than those of PCBs, probably because of the extensive usage of PBDEs in the Philippines. HBCD levels were two orders of magnitude lower than those of PCBs and PBDEs, suggesting that HBCDs are not extensively used in the Philippines. Concentrations of analytes were higher in the eastern part of the bay, implying municipal and industrial areas could be pollution sources of these chemicals. Results of sediment core analysis indicated highest concentrations of PCBs and PBDEs in the sediment layer deposited in the late 1990s. Although no obvious trend was observed for HBCDs due to low levels, it may increase in the future if usage is extended to developing countries.

Keywords: PBDEs, HBCDs, sediment core, Manila Bay

INTRODUCTION

The use of brominated flame retardants (BFRs) has been increasing, especially in the last two decades, and as a result environmental contamination by BFRs is of growing concern (Hites, 2004; Covaci *et al.*, 2006; Law *et al.*, 2008).

Polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecanes (HBCDs) are primarily used as additive BFRs, and are easily released into the environment during use and disposal of the products. Bioaccumulation of PBDEs and HBCDs as well as other persistent organic pollutants such as polychlorinated biphenyls (PCBs) has been reported in quantifiable levels in wildlife (Law *et al.*, 2006; Isobe *et al.*, 2009). Although contamination by BFRs has been a serious problem in Asian developing countries, information on the present status and temporal trend of BFRs pollution is limited. The coastal area is considered to play a major role as a sink for environmental contaminants. Manila Bay, located in the northern Philippine island of Luzon, is one of the largest protected bays of Southeast Asia, with a surface area of 1980 km² and an average water depth of 18 m. About 17 million people live in Metro-Manila and large-scale municipal and industrial activities take place in the watershed of the bay; this condition has caused significant eutrophication to the system. In addition, the bay also receives a wide range of chemicals from agricultural fields, mining areas and an active volcano. Therefore, a survey of anthropogenic contaminants in Manila Bay is necessary to evaluate and monitor the fate and toxicological effects in the tropical aquatic environment. In this study, the spatial distribution and vertical profile of PBDEs and HBCDs as well as PCBs were investigated in 33 surface sediments and a sediment core from Manila Bay, the Philippines.

MATERIALS AND METHODS

Samples

A total of 33 surface sediments and one sediment core were collected from Manila Bay, the Philippines, in June and November 2008. Surface sediment was taken using an Ekman grab and the top 3 cm was sampled. The sediment core was collected using a gravity core sampler and sliced at 2 cm intervals after transporting to the lab. All sediment samples were stored at -25°C in the Environmental Specimen Bank (*es-BANK*) of Ehime University until chemical analysis.

Chemical analysis

Sediment samples were analyzed following the method described elsewhere (Isobe *et al.*, 2007; Minh *et al.*, 2007) with some modifications. Briefly, 5 g of freeze-dried sediment was spiked with surrogates (¹³C₁₂-PBDEs and ¹³C₁₂-HBCDs) and extracted with hexane/acetone using an ultrasonic extractor. The extract was purified with multi-layer column chromatography and concentrated with sulfuric acid. Sulfuric acid was removed by liquid-liquid extraction using hexane and hexane-washed water. The hexane layer was concentrated, cleaned with gel permeation chromatography (GPC), and fractionated with an activated silica gel column chromatography. The fraction containing PBDEs was treated with activated copper strings to remove sulfur. ¹³C₁₂-BDE-139 was spiked as an internal standard and subjected to GC-MS analysis. Concentrations of all the targeted BDE congeners (from mono- to nona-brominated), except BDE-209, were summed




Fig. 1. Concentrations of PCBs, BDE-209, Σ PBDEs and HBCDs in surface sediments from Manila Bay, the Philippines.

to obtain concentration of Σ PBDEs. Concentrations of BDE-209 were shown separately. The HBCDs fraction was evaporated and spiked with HBCDs- d_{18} prior to LC-MS/MS analysis. Concentrations of PBDEs and HBCDs were expressed as ng/g dry weight unless stated otherwise.

RESULTS AND DISCUSSION

Spatial distribution

Organohalogen compounds analyzed in the present study were detected in most of the surface sediment samples, indicating ubiquitous pollution by these contaminants (Fig. 1). Concentrations of PCBs, BDE-209, Σ PBDEs and HBCDs in surface sediments ranged from <0.50 to 110, 0.90 to 110, <0.10 to 18 and <0.02 to 2.8 ng/g dry wt, respectively. Concentrations of BDE-209 were slightly higher than those of PCBs, probably because of the extensive usage of PBDEs in the Philippines. HBCD levels were two orders of magnitude lower than PCBs and PBDEs, suggesting that HBCDs are not extensively used in the Philippines or at least not available to the bay. These levels are baseline data for tracking future use since HBCDs were detected for the first time in this region. Levels of organohalogen compounds in the surface sediments were higher in the eastern part of the bay, in close proximity to the highly populated and industrial area of Metro-Manila, implicating industrial and municipal activities as sources of these compounds. Similar distribution patterns were observed in sediment samples from Tokyo Bay




Fig. 2. Vertical profiles of PCBs, BDE-209, Σ PBDEs and HBCDs in the sediment core from Manila Bay (#1103).

(Minh *et al.*, 2007). Concentrations of BDE-209 were one order of magnitude higher than Σ PBDEs (from mono- to nona-brominated congeners), which is consistent with the other reports dealing with sediment samples.

Vertical profile

Vertical profiles of PCBs, BDE-209, Σ PBDEs and HBCDs in the sediment core (#1103) from Manila Bay are shown in Fig. 2. Results from sediment core analysis indicated increasing trends in the sediment layer from 40 to 10 cm, and a steady state in the upper layers for PCBs and BDE-209. Concentrations of BDE-209 increased from the middle layer (28 cm) while PCBs were detected in the deeper (42 cm) layer. This phenomenon is consistent with the fact that commercial use of deca-BDE mixture started from the 1980s while PCBs had been used since the 1950s. The highest concentrations of PCBs, BDE-209 and Σ PBDEs were observed in the layer deposited in the late 1990s. In recent years Σ PBDEs showed a decreasing trend in the upper layers, which may be a result of the phasing-out of lower brominated PBDE products in the early 2000s. Although no obvious trend was observed for HBCDs due to the low levels, it may increase in the future if usage is extended to developing countries. To prevent future environmental pollution by these persistent contaminants and to evaluate ecological risks, further retrospective studies using sediment cores and archived samples from Asian developing countries are warranted.

Acknowledgments—Financial support was provided by Grants-in-Aid for Scientific Research (S: 20221003) and Global COE Program of the Japanese Ministry of Education, Science, Sports, Culture and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS). This research was also partly supported by MEXT's program "Promotion of Environmental Improvement for Independence of Young Researchers" under the Special Coordination Funds for Promoting Science and Technology and the Global Environment Research Fund (RF-0904) of the Ministry of the Environment.

REFERENCES

- Covaci, A., A. C. Gerecke, R. J. Law, S. Voorspoels, M. Kohler, N. V. Heeb, H. Leslie, C. R. Allchin and J. de Boer (2006): Hexabromocyclododecanes (hbcdds) in the environment and humans: A review. *Environ. Sci. Technol.*, **40**, 3679–3688.
- Hites, R. A. (2004): Polybrominated diphenyl ethers in the environment and in people: A meta-analysis of concentrations. *Environ. Sci. Technol.*, **38**, 945–956.
- Isobe, T., K. Ramu, N. Kajiwara, S. Takahashi, P. K. Lam, T. A. Jefferson, K. Zhou and S. Tanabe (2007): Isomer specific determination of hexabromocyclododecanes (hbcdds) in small cetaceans from the south china sea-levels and temporal variation. *Mar. Pollut. Bull.*, **54**, 1139–1145.
- Isobe, T., Y. Ochi, K. Ramu, T. Yamamoto, Y. Tajima, T. K. Yamada, M. Amano, N. Miyazaki, S. Takahashi and S. Tanabe (2009): Organohalogen contaminants in striped dolphins (*Stenella coeruleoalba*) from Japan: Present contamination status, body distribution and temporal trends (1978–2003). *Mar. Pollut. Bull.*, **58**, 396–401.
- Law, K., T. Halldorson, R. Danell, G. Stern, S. Gewurtz, M. Alaee, C. Marvin, M. Whittle and G. Tomy (2006): Bioaccumulation and trophic transfer of some brominated flame retardants in a Lake Winnipeg (Canada) food web. *Environ. Toxicol. Chem.*, **25**, 2177–2186.
- Law, R. J., D. Herzke, S. Harrad, S. Morris, P. Bersuder and C. R. Allchin (2008): Levels and trends of hbcdd and bdes in the European and Asian environments, with some information for other bfrs. *Chemosphere*, **73**, 223–241.
- Minh, N. H., T. Isobe, D. Ueno, K. Matsumoto, M. Mine, N. Kajiwara, S. Takahashi and S. Tanabe (2007): Spatial distribution and vertical profile of polybrominated diphenyl ethers and hexabromocyclododecanes in sediment core from Tokyo bay, Japan. *Environ. Pollut.*, **148**, 409–417.

T. Isobe (e-mail: t.isobe@agr.ehime-u.ac.jp)