

Age-validated Longevity of Fishes: Its Importance for Sustainable Fisheries

Gregor M. Cailliet*
and
Allen H. Andrews

*Moss Landing Marine Laboratories
8272 Moss Landing Road
Moss Landing, CA 95039, USA*

*E-mail: Cailliet@mml.calstate.edu

Marine fisheries have spread to the deep-sea because the fishes in many of the shallow-water marine habitats have been overexploited. Many studies have found that deep-sea fishes generally grow slowly and can achieve long life spans; however, most age estimates have not been validated. Several methods for validating age and growth estimates have been developed and applied to deep-water and other long-lived organisms. These include lead-radium and bomb radiocarbon dating of calcified structures. Application of these techniques has usually validated high lifespan estimates, some in excess of a century. In this paper, we discuss the applications and limitations of these techniques for difficult-to-age fishes. Because many fishery management techniques focus on biomass or yield and not as much on life history stages of fished populations, we specifically focus on the relevance of age validation to fishery management of deep-dwelling fishes, especially relative to slow growth, late age at maturity, long life spans, and the relative contribution of big, old, fecund females (BOFFs) and maternal-age-dependent larval survival to future generations via lifetime fecundity. These life history traits make their populations more vulnerable to fishing mortality, thus emphasizing the need to accurately assess longevity in fishes.

KEYWORDS age validation; bomb radiocarbon; carbon-14 ($\Delta^{14}\text{C}$); deep-water fishes; fecundity; fishery management; growth; life history; life span; lead-radium dating; longevity; maternal-age-dependent larval survival; mortality; vulnerability

1. Introduction

Many marine fisheries are reported as overexploited on a global scale (Pauly *et al.* 2003). This overexploitation appears to have influenced the relative abundance of predatory fishes in some marine ecosystems (Pauly 1995; Worm *et al.* 2005) and may have upset the balance of some major ecosystems (Pauly and Christensen 1995; Pauly *et al.* 1998, 2002, 2003; Christensen *et al.* 2003; Frank *et al.* 2005). There is argument about the extent and the causes of these fishery declines and ecological impacts (Pauly 1995; Pauly *et al.* 1998, 2002, 2003; Myers *et al.* 2003; Myers and Worm 2003, 2005; Walters 2003; Hutchings and Reynolds 2004; Safina *et al.* 2005; Sibert *et al.* 2006; Worm *et al.* 2005; Hilborn 2007). However, it is certain that many near-shore fisheries have been so highly overexploited that fishing effort has moved further offshore and into deeper water (Clark 2001; Gordon 2001; Haedrich *et al.* 2001; Roberts 2002; Stockley *et al.* 2005).

Fishery modeling and management practices have primarily focused on stock assessments, which mainly use biomass and maximum (MSY) or optimum (OSY) sustainable yield, to attempt to determine the level of exploitation or sustainability of a given fishery (Hilborn and Walters 1992; Gallucci *et al.* 1996; Berkeley 2006). In some cases, numerical abundance and/or age composition were taken into consideration, but even age-structured analyses often make major assumptions about age-driven biological quantities. Because the complexities of population age-structure are poorly understood, dynamic parameters that are an important consideration must be modeled or assumed to be fixed (e.g., Methot 1990; Millar and Methot 2002). Stock assessments are now commonly performed, but the concept of sustainability is still very much in question (Millar and Methot 2002; Methot and Piner 2002; Ralston *et al.* 2003; Beamish *et al.* 2006; Longhurst 2006).

Fishing practices in many cases have caused the size (and presumably age) structure of many populations to be fished down (Mason 1998; Pauly *et al.* 1998; Worm *et al.* 2005; Harvey *et al.* 2006), and this may have seriously impacted long-lived fishes like chondrichthyans and those in the deep sea. In nearshore environments along the west coast of the U.S. there is evidence of significant shifts downward in the size composition of rockfish landings (Mason 1998). This fished-down size and age structure can have numerous impacts and a few of the most significant are further discussed here.

First, it can cause a loss of genetic diversity or heterozygosity (e.g., Hauser *et al.* 2002). In addition, some fishes are genetically structured at the regional and sub-regional scales, as was demonstrated for Patagonian toothfish (*Dissostichus eleginoides*; Rogers *et al.* 2006) and further described in otolith elemental analyses (Ashford *et al.* 2007), which influences how fishing might affect the genetic structure of populations. Stockley *et al.* (2005) demonstrated that the deep-sea black-spot sea bream (*Pagellus bogaraveo*), which was heavily fished in recent years due to depletion of shallow-water fish stocks, may have undergone a severe bottleneck in its genetic population structure; low genetic variability in its populations was documented due to the small population size that resulted from fishing.

Second, removing the larger individuals can favor slower-growing and earlier-maturing fishes (Kenchington 2002; Ziegler *et al.* 2007). Thus heritable differences in yield and life-history traits can be caused by selective fishing which has been demonstrated for fishes such as the northern cod (*Gadus morhua*; Novikov *et al.* 2002; Hutchings 2004; Olsen *et al.* 2004).

Third, there can also be long term ecological and evolutionary implications of size-specific fishing mortality, resulting in a "dynamic tug-of-war between natural and harvest selection" (Edeline *et al.* 2007) in

terms of size and age at first reproduction and size- and age-specific reproductive output (Conover and Munch 2002; Berkeley *et al.* 2004a; Bobko and Berkeley 2004; Berkeley 2006; de Roos *et al.* 2006; Walsh *et al.* 2006; Conover 2007; Jorgensen *et al.* 2007).

Fourth, there can be ecosystem effects (Frank *et al.* 2005; Scheffer *et al.* 2005; Worm *et al.* 2005) that can be manifested as loss of top carnivores and changes in trophic relationships of ecosystems. One management technique that can help alleviate this problem, at least on a local scale, is the implementation of Marine Protected Areas (MPAs; National Research Council 2001). These are being seriously considered and implemented in several places around the world, especially along the west coast of the United States (Parker *et al.* 2000). Berkeley (2006), however, pointed out that MPAs do not seriously consider the life histories of fished populations, except as sources of larvae for export from mature females.

Fifth, size-selective fishing has been estimated to cause a subsequent reduction in lifetime production, measured by O'Farrell and Botsford (2005, 2006a, b) as lifetime egg production (LEP) in rockfishes (Genus *Sebastes*; Family Scorpaenidae) and by Harvey (2005) as age-specific egg production. In addition, it can reduce survivorship of larvae from "fit" or well-adapted adults. It has been shown for some species that the larvae from these older females in some rockfishes are more numerous and robust, being stronger and more likely to survive (Love *et al.* 2002; Berkeley *et al.* 2004a, b; Bobko and Berkeley 2004; Berkeley 2006). And those larger, most likely older, females not only are often more fecund but they can also exhibit maternal-age-dependent increased larval survival. This has caused some authors to contend that fisheries, in general, need to "leave the big ones" (Birkeland and Dayton 2005).

2. The Importance of Age Validation in Estimating Longevity (Lifespan)

We propose here that the life history parameters longevity (length of life) and/or lifespan (average time between birth and death) are very important to consider when determining how to manage exploited populations. Long-lived fishes have often had their ages and life spans underestimated, causing fishery management policies to be less effective. This is not a new concept because the importance of age validation was strongly emphasized more than two decades ago by Beamish and MacFarlane (1983) and more recently by Campana (2001), among many other authors. However, many age, growth and longevity studies have not provided validated age estimates leading to the parameters determined in their studies.

Traditional age validation methods (mark-and recapture, laboratory rearing, or marginal increment analysis) are often difficult or impractical to use for long-lived or deep-sea fishes (Campana 2001). There was little hope for such applications to deepwater fishes until recent developments provided some promise for tagging deep-sea species *in situ*. This was pioneered by Starr *et al.* (2002) for eastern Pacific deep-water rockfishes, and used in other deep-sea applications (e.g. Sigurdsson *et al.* 2006) in deep water (~500–800 m) off Iceland on redfish (*Sebastes mentella*).

In lieu of such applications, a naturally occurring radioisotope pair found in calcified structures of fishes, radium-226 (^{226}Ra) and its daughter product lead-210 (^{210}Pb), can act as a built-in chronometer. In addition to this radiometric technique, atmospheric testing of thermonuclear devices in the 1950s and 1960s caused a rise in radiocarbon ($\Delta^{14}\text{C}$) and produced a globally distributed signal that diffused into the ocean surface worldwide (Broecker and Peng 1982).

In recent years, many of the age estimates for deep sea fishes have been reevaluated and

in many cases were deemed drastically greater than previously suspected. A number of these age estimates, ones that were often met with disbelief because of the great longevity estimates, have been validated through use of lead-radium and bomb radiocarbon dating, some of which also included tag-recapture and oxytetracycline from field studies (e.g., Beamish and McFarlane 1983; Kestelle *et al.* 1994). Application of these radiometric methods has been successful for deep-sea fishes (e.g. Andrews *et al.* 1999, 2007) and this is especially true for a numerous rockfishes (Andrews *et al.* 2002, 2005, 2007; Bennett *et al.* 1982; Campana *et al.* 1990; Kestelle *et al.* 2000; Kestelle and Kimura 2006; Kerr *et al.* 2004, 2005; Piner *et al.* 2005; Stevens *et al.* 2004; Stransky *et al.* 2005; Watters *et al.* 2006). In some cases, complications in what appears to be a system that was not conserved with time provided circumstances that violated the necessary assumptions for successful application (e.g., cartilage of chondrichthyans and sturgeon; Burton *et al.* 1999; Welden *et al.* 1987). For otolith material, however, numerous other bony fishes have been validated using lead-radium and bomb radiocarbon dating (e.g., Fenton *et al.* 1991; Smith *et al.* 1991, 1995; Campana 1997; Andrews *et al.* 2001; Kalish *et al.* 2001; Ewing *et al.* 2007).

In this paper, we return to Beamish and McFarlane's (1983) plea for age validation to become a more common element in fish age and growth studies. In doing so, we will demonstrate the importance of age-validated longevity to fish population dynamics, especially in exploited conditions. We will also incorporate the results and concepts initiated by Berkeley *et al.* (2004a, b), Bobko and Berkeley (2004), and Berkeley (2006), in which they demonstrated the importance of larger, fecund female rockfishes (recently coined as "Big Old Fecund Females" or BOFFs), using the black rockfish (*Sebastes melanops*) aged 50 years old by counting rings in sectioned otoliths.

O'Farrell and Botsford (2005, 2006a, b) used size structure of the same species (*S. melanops*), but also included five other rockfish species to demonstrate the effect of size-selective fishing on LEP. They assumed that size classes represented age classes, but no age validation has been accomplished for these species. Similar studies have been done on rockfishes by Harvey (2005) and Harvey *et al.* (2006), assuming specific growth characteristics, not all of which were validated. Thus, it is important to note that the ages estimated or inferred from size-composition or otolith readings could leave some doubt about the reality of the BOFF and/or maternal-age-dependent larval survival, and perhaps the reduction in LEP. Hence, we emphasize use of age-validated species and that this perspective on age validation becomes more important when taking the previously stated factors into consideration.

3. The Importance of Lifetime Fecundity (Reproductive Output)

It is important to note that true reproductive output is a factor that can only be obtained by validating longevity or lifetime of a given species (Harvey 2005; O'Farrell and Botsford 2006a, b). In this paper, two terms will be used with regard to reproduction; fertility, the actual reproductive productivity or production of offspring, and fecundity, the physiological potential for reproduction or the age-specific state of being fertile. It appears that most fishes, including rockfishes, continue to reproduce after maturity, without evidence of senescence in all years following the age at first maturity (Cailliet *et al.* 2001). Because larger, and presumably older, fish often have higher fecundity and may have better adapted larvae, their lifetime fecundity becomes an important parameter in understanding and predicting population dynamics (Phillips 1964; Gunderson *et al.* 1980; Haldorson and Love 1991; Lea *et al.* 1999; Love *et al.* 1990, 2002; Berkeley

et al. 2004a, b; Bobko and Berkeley 2004; Harvey 2005; Berkeley 2006).

Summarized data on size-specific fecundity for rockfishes demonstrated that larger rockfishes typically produce more offspring (Love *et al.* 1990, 2002). Thus, fishing these larger size and older age classes heavily can truncate the number of reproductive years and thus reduce their ability to maintain viable populations. This has been demonstrated demographically by Ralston *et al.* (2003) for the shortbelly rockfish (*Sebastes jordani*) and by Ziegler *et al.* (2007) on the banded morwong (*Cheilodactylus spectabilis*). Also, environmental processes, like El Niño events, can also have a serious influence on reproductive output like egg production in fishes (Ven Tresca *et al.* 1995; Harvey 2005). We propose that this approach, using age-validated lifetime fecundity, should be a major research objective for all exploited fishes, including rockfishes.

Demographic approaches using either life table analyses or stage- and/or age-based matrices are good ways to produce predicted population dynamics, with and without exploitation (Cailliet 1992; Ebert 1999; Caswell 2001; Mollet and Cailliet 2002, 2003). In addition, sensitivity and elasticity analyses (Heppell *et al.* 2000; Frisk *et al.* 2005; Garcia *et al.* 2008) will allow the most ecologically important life stages to be identified and used in fishery management policy. However, it may be that sustainability is unachievable (Longhurst 2002; 2006), especially without knowing these age-specific life history traits.

4. Deep-water Rockfish Age Determination, Validation, and Longevity

For this paper, we have chosen a few examples of age validation research on rockfish populations with which we are familiar off the west coast of the United States (Andrews *et al.* 2002, 2005, 2007; Kerr *et al.* 2004; Piner *et al.* 2005). We will focus on three

species of rockfishes for which age has been validated, and life spans determined based on the maximum growth zone counts, for which one or both of the radiometric techniques (lead-radium or bomb radiocarbon dating) was used to support estimates of age.

Our bomb radiocarbon age-validation results on bocaccio (Andrews *et al.* 2005; *Sebastes paucispinis*) are useful relative to the papers cited above on black rockfish (*S. melanops*) because they are similar in terms of lifespan (~50 years), but perhaps not with regard to the other parameters as studied by Berkeley *et al.* (2004a, b) and Bobko and Berkeley (2004). We investigate both their size and age at maturity and their fecundity in this regard. We also present age-validated results for two other species on which we have published age-validation papers, the canary (*Sebastes pinniger*) and yelloweye (*Sebastes ruberrimus*) rockfishes, both of which have had their ages (and life spans) validated using lead-radium and bomb radiocarbon dating (Andrews *et al.* 2002, 2007; Kerr *et al.* 2004; Piner *et al.* 2005).

For these and other age estimation and validation studies, calcified structures such as otoliths, vertebrae, other bones, spines, and thorns have often proved useful and accurate. These structures are prepared for age determination in ways that enhance the readability of growth zones that are typically present. In age validation studies the use of core or growth zone specific material from adult structures and from various year classes have provided better age resolution than previously possible for both lead-radium and bomb radiocarbon dating.

Successful application of lead-radium and bomb radiocarbon dating to rockfishes (family Scorpaenidae) has a well developed history. The first application of lead-radium dating to otoliths in general was an age validation study on splitnose rockfish (*S. diploproa*) in which a 60 yr lifespan was supported from the growth-zone derived age estimation criteria (Bennett *et al.* 1982).




Fig. 1. Diagrammatic representation of the theory behind lead-radium dating. Represented is the ingrowth of lead-210 from radium-226 as the ingrowth curve in the figure. The clusters of hypothetical data points represent the various forms of potential age differentiation from lead-radium ratios measured from samples with estimated age. The clusters not in agreement with the ingrowth curve represent scenarios where age was either underestimated or overestimated by several decades. The margin of error for age is typical for an age distribution from a pooled group of otoliths. The margin of error for lead-radium ratios for lead-radium activities that would be somewhat ideal. See Fig. 2 for a different graphic analysis of these same theoretical data examples.

Since this pioneering application, the technique has been applied to several other rockfishes with varying degrees of age resolution (Burton *et al.* 1999; Campana *et al.* 1990; Kestelle *et al.* 2000; Andrews *et al.* 2002, 2005, 2007; Stevens *et al.* 2004; Watters *et al.* 2006). Bomb radiocarbon dating of fish otoliths began in the southern hemisphere (Kalish 1993, 2001), followed by a series of studies in the northern hemisphere (e.g., Campana 1997; Kalish *et al.* 2001).

Lead-radium dating relies on the incorporation of naturally occurring radium-226 from the environment into the otolith and its subsequent decay to lead-210. By measuring the disequilibria of these two radioiso-

topes in otolith core material (first few years of life), an independent estimate of age can be determined based on the known ingrowth rate of lead-210 from radium-226 (Fig. 1; Campana *et al.* 1990; Smith *et al.* 1991; Kimura and Kestelle 1995; Francis 2003). This technique works well as a tool for determining the validity of age interpretations that differ considerably, but its application is limited by relatively low resolution at ages approaching 100 years. Typically, the end result for this kind of study is support for, or refutation of, a given age estimation technique and establishment of an independent estimate of age or lifespan. For example, lead-radium dating performed on the yelloweye rockfish (*S. ruberrimus*) provided




Fig. 2. Comparison of otolith-generated age estimate to radiometric age estimate using ratio of lead-210 to radium-226. The cluster of value on the line indicate that age estimates were confirmed by lead-radium age estimates, while those to the upper left mean that lead-radium ages were higher than growth zone derived ages. At the lower right the values indicate that otolith-derived ages were too high and lead-radium age estimates indicated that these samples were younger. See Fig. 1 for a different graphic analysis of these same theoretical data examples.

support for age exceeding ~100 yr (Andrews *et al.* 2002). The application refuted age estimates that were significantly younger and provided age validated support for transverse otolith ageing techniques, but this finding was the extent of the maximum age resolution because of technique limitations. These findings were similar for bocaccio rockfish for which a minimum age of 31 years was determined using lead-radium dating (Andrews *et al.* 2005).

Many papers now have been written on radiometric age determination of fishes since Bennett *et al.* (1982) first confirmed the longevity of the splitnose rockfish (*Sebastes diploproa*) using lead-radium dating. Such age validation studies usually involve two

graphical comparison techniques. The first involves plotting the predicted ingrowth curve over time, plotted with individual age estimates (usually from growth zone counts in otolith sections), to determine the agreement between measured and expected lead-radium ratios (Fig. 1). The second approach is to plot the growth zone derived age estimates versus the radiometric age estimate for each sample (Fig. 2). With this kind of comparison, it is possible to test for trends of agreement or disagreement based on the hypothesis that there is a one-one agreement that results from a regression fitted to the data. The one discrepancy in this process that is still a work in progress is taking into consideration the change of radiometric age uncertainty with increasing age.

Use of the bomb radiocarbon chronometer requires a series of individual otoliths for which the birth years, based on standardized growth-zone age estimates, range from a time prior to significant atmospheric testing of thermonuclear devices (pre-1957) to the post-bomb period. This approach can utilize both the 1) time of first rise in radiocarbon, and the 2) sloped period for the rise in radiocarbon, as a time-reference markers for age validation. It is the agreement of the bomb radiocarbon record, from the species with age in question, with a reference bomb radiocarbon time-series that provides a basis for age validation. Hence, the utility of this approach for determining age or lifespan is dependent upon the difference between the collection year and time of first rise in radiocarbon for the reference time series. For example, a minimum lifespan of 43 ± 3 years was determined for canary rockfish (*Sebastes pinniger*) based on the first sample to have measured radiocarbon levels that were pre-bomb (Andrews *et al.* 2007). In addition, age can be determined by projecting measured radiocarbon levels back to a reference time series (Campana 1997); a minimum life span of 37 ± 2 years was determined in this manner for bocaccio rockfish (Andrews *et al.* 2005). A thorough evaluation of age classes can also be undertaken with a series of samples with different collection years (Piner and Wischniowski 2004).

For each of the three species discussed here, age has been determined using both lead-radium and bomb radiocarbon dating. Lead-radium results for yelloweye rockfish provided strong support for the age estimation criteria used to determine ages that exceed 100 years and the consistency of fit to the ingrowth curve for younger age classes provided additional support. Based on the ease of otolith reading for this species using the same criteria, we considered a radiocarbon record from otoliths of yelloweye rockfish would provide a reference chronology for validation of other regional fish spe-

cies (e.g., Kerr *et al.* 2005). Some (e.g. Stewart and Piner 2007) have questioned this assumption and suggested that the follow up study using radiocarbon was instead further validation for the age estimate criteria and further support for the findings of the lead-radium study.

It seems there is an element of truth in both perspectives. Based on the findings of the combined studies, and depending on your perspective, the evidence that yelloweye rockfish is one of the longest lived rockfishes is robust. Lead-radium results for bocaccio rockfish provided only a minimum age for this species because of the remarkably low radium-226 levels; there was 95% confidence that the minimum age this species can attain is 31 years. This finding was well supported and exceeded with the application of bomb radiocarbon dating. Lead-radium and bomb radiocarbon results for canary rockfish worked well together by providing robust support for ages estimated in the Canadian canary rockfish fishery. Each application has had variable degrees of age validation based on the uncertainty or assumptions of the application, and there is still a need for developing a high-precision dating method.

Here, we stress the paramount importance that some form of validation or support be given for age estimation procedures, and ultimately estimates of longevity, before we can accurately understand the population dynamics of a population supporting a given fishery. Without determining the temporal periodicity of the growth zones in their otoliths, the variation in growth and longevity would not be fully understood for different species of fishes like bocaccio, canary and yelloweye rockfishes.

5. Deep-water Rockfish Age-Specific and Lifetime Reproductive Output

For the three species of interest here, we have provided age validated support using both lead-radium bomb radiocarbon dating (e.g.,




Fig. 3. Plot of measured bomb radiocarbon ($\Delta^{14}\text{C}$) levels in otolith cores, representing the first year of growth, from adult bocaccio (*S. paucispinis*; Andrews *et al.* 2005), canary (*Sebastes pinniger*; Andrews *et al.* 2007), and yelloweye (*S. ruberrimus*; Kerr *et al.* 2005) rockfishes, plotted with $\Delta^{14}\text{C}$ data from whole otoliths of juvenile Pacific halibut (*Hippoglossus stenolepis*; Piner and Wischniowski 2004). Birth year was estimated using growth zone counts for both canary and bocaccio rockfishes and measured $\Delta^{14}\text{C}$ levels from each aged core was compared to the chronological reference time series established for the Northeastern Pacific Ocean; the chronological time series was established for the region with the age-validated yelloweye rockfish (Andrews *et al.* 2002) and known age juvenile Pacific halibut otolith records. Error bars for the data represent the analytical uncertainty of the $\Delta^{14}\text{C}$ measurements and the potential range for the birth date based on the age estimate uncertainty.

Fig. 3). For bocaccio, canary, and yelloweye rockfishes, the age estimates from otolith growth zone counts produced age estimates that conformed well to known regional bomb radiocarbon levels. Thus, age estimation procedures are independently supported for the estimated life spans of these three fishes and it is this supported life span that we propose is an important factor in determining the sustainability of fisheries.

This allows us to also feel confident that the resulting growth curves (Fig. 4 from Love *et al.* 2002) provide evidence of a variety of growth characteristics and life spans among

many species of the genus *Sebastes*. Growth curves for 13 species of rockfishes, including yelloweye and canary rockfishes, are plotted in Fig. 4, but no growth curve was yet available for bocaccio when Love *et al.* (2002) published this figure. We now have a better idea of bocaccio growth, with best estimates, taken from MacCall (2007), of growth coefficients (K) ranging from 0.184–0.210, and asymptotic lengths (L_{∞}) being 75.9 and 65.6 cm TL, for females and males respectively. A life span estimate of ~50 years was used by Love *et al.* (2002). However, there is considerable variation in the value




Fig. 4. A compilation of data from numerous studies where von Bertalanffy growth curves were fitted to female length-at-age data (otolith section age) for 13 rockfish species. In the context of this study, the bocaccio growth data were not plotted because at the time of publication, age data was not yet resolved (Data and figure from Figure 9.6, page 61, in Love *et al.* (2002) with permission to reprint).

used for maximum age. MacCall (2007) reduced his maximum age estimate to 27 from 45 years used in his original 2003 stock assessment. Our most recent work indicates 37 ± 2 years for the largest fish used in the study (Andrews *et al.* 2005). Therefore, we feel the best estimate would be ~ 37 years based on our age-validated results, the importance of which will be discussed further.

A synopsis (Love *et al.* 2002) of fecundity information gathered from numerous studies has produced evidence that larger, and presumably older, rockfish females can provide maximum fecundities (number of eggs or number of larvae produced) that increase exponentially with maximum length (Fig. 5). Thus, the deeper-dwelling species, which are often larger, may have increased fecundity coupled with their increased longevity (Cailliet *et al.* 2001). The important question that comes from this observation

has to do with the number of years past age at maturity over which these females continue to produce eggs and larvae. In addition, both fishing and natural mortality can cause the relative numbers of females in successively older age classes to decline, further reducing their net reproductive rates over these years.

Knowing the longevity and age-specific fecundity for rockfishes allows us to speculate about the potential impact of size-specific mortality from fishing. Without fishing mortality, bocaccio, canary and yelloweye rockfishes would start spawning from the ages of approximately 7, 8, and 19 years, to the end of their lifetime at ages approaching 37, 84, and 118 years, respectively (Berkeley 2006; modified by our recent results). Hence, the longer-lived, deeper-dwelling canary and yelloweye rockfishes would have up to 76 and 99 reproductive years, versus up to 30




Fig. 5. A compilation of data used to establish a relationship between rockfish maximum length (L , cm) and fecundity (F) for 33 rockfish species from the eastern North Pacific (Data and figure from Figure 5.8, page 37, in Love *et al.* (2002) with permission to reprint). This relationship may be an indication that larger, and presumably older, fish are typically more fecund; typifying the concerns associated with size-specific management strategies.

years for the shallower, shorter-lived bocaccio rockfish. By taking this into consideration, it is relatively straight forward to predict that size-specific fishing mortality (i.e. heavily fishing the larger, older fish) would have a deleterious impact on lifetime egg (larval) production. This would be even more convincing if age-specific natural mortality, the growth rate and age at maturity were known.

Using this line of thinking, it is interesting to consider several other rockfishes that have greater life spans. The rougheye (*Sebastes aleutianus*) and shorttraker (*Sebastes borealis*) rockfishes have estimated longevitys ranging up to 157 and 205 years, respectively (Munk 2001); however, these estimates are based on growth zone counts from otolith sections. Neither lead-radium or bomb radiocarbon dating has been applied and the development of other tech-

niques is necessary to accurately ascertain ages of this magnitude. Assuming these estimates are accurate, these exceedingly long-lived fishes would be more vulnerable to size-specific fishing mortality that targets deep-dwelling individuals that are larger and potentially much older.

Likewise, age validation is an important factor in managing other long-lived fishes that are heavily exploited. For example, Baker and Wilson (2001) and Milton *et al.* (1995) validated longevitys in snappers (family Lutjanidae) occupying relatively deep water in the Gulf of Mexico and off Australia, respectively. Another prime example, with a bit of a twist, is the orange roughy (*Hoplostethus atlanticus*) with its contrasting reproductive strategy of bestowal (maximizing investment in a smaller number of offspring; Pankhurst and Conroy 1987). This factor, combined with its validated

centenarian longevity (Fenton *et al.* 1991; Smith *et al.* 1995; Andrews and Tracey 2007), make orange roughly a particular concern in a size-selective management strategy. Other deep-water fishes with potentially high longevity, like oreos (family Oreosomatidae: Stewart *et al.* 1995), need to be further considered in this regard. This kind of vulnerability would be especially true for those targeted preferentially for egg production, like anglerfishes (family Lophiidae, genus *Lophius*: Duarte *et al.* 2001; Landa *et al.* 2001).

6. Conclusions

It is a combination of age structure (keep the big ones), life-time egg production (greater productivity), and genetic diversity (potential enhanced survivorship) that beg for further consideration of age validation and abandonment of only size-specific management practices. Age validation for fishes that are recognized as long-lived (e.g., Burton *et al.* 1999 and Cailliet *et al.* 2001), such as chondrichthyans (sharks: Cailliet and Goldman 2004; Cailliet *et al.* 2006), chondrosteans (sturgeon & paddlefishes: Burton *et al.* 1999), and tarpons (Andrews *et al.* 2001), either have required or will need to use a combination of techniques, including bomb radiocarbon, lead-radium dating, and/or tag-recapture and marking techniques to validate their growth characteristics and longevity estimates.

For chondrichthyan fishes and sturgeons, which are cartilaginous and do not have calcified otoliths that are useful, lead-radium dating did not work due to what appeared to be a violation of the closed system assumption (Welden *et al.* 1987; Burton *et al.* 1999). Radionuclides may move around from the vertebral column to the rest of the body, thus making lead-radium dating not useful for these fishes.

For some elasmobranch species, however, bomb radiocarbon dating has been suc-

cessfully applied because the signal is conserved in cartilage. Apparently, the bomb radiocarbon signal is conserved for some species, but may be: 1) free to move around in the vertebral column; or 2) mixed with radiocarbon-depleted sources in other species. This technique has worked well for the porbeagle (*Lamna nasus*: Campana *et al.* 2002), shortfin mako (*Isurus oxyrinchus*: Campana *et al.* 2002; Ardizzone *et al.* 2006), spiny dogfish (*Squalus acanthias*: Campana *et al.* 2006), and tiger shark (*Galeocerdo cuvieri*: Kneebone *et al.* in press). It did not work as well for white shark (*Carcharodon carcharias*: Kerr *et al.* 2006).

In summary, both lead-radium and bomb radiocarbon dating provide solid, age-based evidence as tools in age validation of fishes. Lead-radium dating is useful for the groups of fishes with otoliths large enough, and with enough radioactivity, for core applications. Bomb radiocarbon dating has often corroborated age estimates, growth characteristics, and life spans of rockfishes, and deeper-dwelling species. Results so far indicate that these deep-water fishes can live a long time, and in a growing number of cases exceed a centenarian life span. As pointed out by Longhurst (2002, 2006), longevity is extremely important to insure effective, age-based management practices. In addition, the metabolism and growth of deep-water fishes appear to be lower and slower than near-shore fishes (Haedrich *et al.* 2001; Roberts 2002; Marriott *et al.* 2006; Drazen and Seibel 2007). Prevention of fishing down the larger and more productive age, not just size, classes is of paramount importance.

Acknowledgments

We thank Bob Lauth (Northwest Fisheries Science Center, National Marine Fisheries Service [NMFS], National Oceanic and Atmospheric Administration [NOAA]); and Craig Lundstrom (University of Illinois, Urbana-Champaign). This paper was funded in part by the National Sea Grant College Program, the National Oceanic and Atmospheric Administration,

U.S. Department of Commerce, under grant numbers NA36RG0537, NA06RG0142, and NA06RG0477, project numbers R/F-148, R/F-174, R/F-176, R/F-182, R/F-190, through the California Sea Grant College System, and in part by the California State Resources Agency. The views expressed herein are those of the authors and do not necessarily reflect the views of NOAA or any of its sub-agencies. The U.S. government is authorized to reproduce and distribute for governmental purposes. We were also supported, in part, under the auspices of the U.S. Department of Energy, by the University of California, Lawrence Livermore National Laboratory under Contract No. W-7405-Eng-48.

References

- Andrews AH, Tracey DM. Age validation of orange roughy and black cardinalfish using lead-radium dating. Final Research Report for Ministry of Fisheries, Res. Project DEE2005/02. NIWA, Wellington, New Zealand. 2007; 40 pp.
- Andrews AH, Cailliet GM, Coale KH. Age and growth of the Pacific grenadier (*Coryphaenoides acrolepis*) with age estimate validation using an improved radiometric ageing technique. *Can. J. Fish. Aquat. Sci.* 1999; **56**: 1339–1350.
- Andrews AH, Burton EJ, Coale KH, Cailliet GM, Crabtree RE. Application of radiometric age determination to the Atlantic tarpon, *Megalops atlanticus*. *Fishery Bull.* 2001; **99**: 389–398.
- Andrews AH, Cailliet GM, Coale KH, Munk KM, Mahoney MM, O'Connell VM. Radiometric age validation of the yelloweye rockfish (*Sebastes ruberrimus*) from southeastern Alaska. *Mar. Freshwat. Res.* 2002; **53**(2): 1–8.
- Andrews AH, Burton EJ, Kerr LA, Cailliet GM, Coale KH, Lundstrom CC, Brown TA. Bomb radiocarbon and lead-radium disequilibria in otoliths of bocaccio rockfish (*Sebastes paucispinis*): a determination of age and longevity for a difficult-to-age fish. *Mar. Freshwat. Res.* 2005; **56**: 517–528.
- Andrews AH, Kerr LA, Cailliet GM, Brown TA, Lundstrom CC, Stanley RD. Age validation of canary rockfish (*Sebastes pinniger*) using two independent otolith techniques: lead-radium and bomb radiocarbon dating. *Mar. Freshwat. Res.* 2007; **58**: 531–541.
- Ardizzone D, Cailliet GM, Natanson LJ, Andrews AH, Kerr LA, Brown TA. Application of bomb radiocarbon chronologies to shortfin mako (*Isurus oxyrinchus*) age validation. *Env. Biol. Fishes* 2006; **77**: 355–366.
- Ashford JR, Jones CM, Hofmann EE, Everson I, Moreno CA, Duhamel G, Williams R. Otolith microchemistry indicates population structuring by the Antarctic Circumpolar Current. *Can. J. Fish. Aquat. Sci.* 2007; **65**: 135–146.
- Baker MS, Wilson CA. Use of bomb radiocarbon to validate otolith section ages of red snapper *Lutjanus campechanus* from the northern Gulf of Mexico. *Limnol. Oceanogr.* 2001; **46**: 1819–1824.
- Beamish RJ, McFarlane GA. The forgotten requirement for age validation in fisheries biology. *Trans. Am. Fish. Soc.* 1983; **112**: 733–743.
- Beamish RJ, McFarlane GA, Benson GA. Longevity overfishing. *Prog. Oceanogr.* 2006; **68**: 289–302.
- Bennett JT, Boehlert GW, Turekian KK. Confirmation of longevity in *Sebastes diploproa* (Pisces: Scorpaenidae) from $^{210}\text{Pb}/^{226}\text{Ra}$ measurements in otoliths. *Mar. Biol.* 1982; **71**: 209–215.
- Berkeley SA. Pacific rockfish management: are we circling the wagons around the wrong paradigm? *Bull. Mar. Sci.* 2006; **78**(3): 655–668.
- Berkeley SA, Chapman C, Sogard SM. Maternal age as a determinant of larval growth and survival in a marine fish, *Sebastes melanops*. *Ecology* 2004a; **85**(5): 1258–1264.
- Berkeley SA, Hixon MA, Larson RJ, Love MS. Fisheries sustainability via protection of age structure and spatial distribution of fish populations. *Fisheries* 2004b; **29**(8): 23–32.
- Birkeland C, Dayton PK. The importance in fishery management of leaving the big ones. *Trends Ecol. Evol.* 2005; **20**(7): 1–3.
- Bobko SJ, Berkeley SA. Maturity, ovarian cycle, fecundity, and age-specific parturition of black rockfishes (*Sebastes melanops*). *Fishery Bull.* 2004; **102**: 418–429.
- Broecker WS, Peng TH. The anthropogenic invasion: the movement of water through the oceanic thermocline. *In: Tracers in the Sea*. Lamont-Doherty Geological Observatory, New York. 1982; 383–444.
- Burton EJ, Andrews AH, Coale KH, Cailliet GM. Application of radiometric age determination to three long-lived fishes using $^{210}\text{Pb}/^{226}\text{Ra}$ disequilibria in calcified structures: a review. *In: Musick JA (ed). Ecology and Conservation of Long-Lived Marine Animals*. Am. Fish. Soc. Spec. Pub. 1999; **23**: 77–87.

- Cailliet GM. Demography of the central California population of leopard shark (*Triakis semifasciata*). *Aust. J. Mar. Freshwat. Res.* 1992; **43**(1): 183–193.
- Cailliet GM, Goldman KH. Age Determination and Validation in Chondrichthyan Fishes. In: Carrier J, Musick JA, Heithaus MR (eds). *Biology of Sharks and Their Relatives*. CRC Press, Boca Raton, FL. 2004; Chapter 14, pp. 399–447.
- Cailliet GM, Andrews AH, Burton EJ, Watters DL, Kline DE, Ferry-Graham LF. Age determination and validation studies of marine fishes: do deep-dwellers live longer? *Exp. Ger.* 2001; **36**: 739–764.
- Cailliet GM, Smith WD, Mollet HF, Goldman KJ. Age and growth studies of chondrichthyan fishes: the need for consistency in terminology, verification, validation, and growth function fitting. *Env. Biol. Fishes* 2006; **77**: 211–228.
- Campana SE. Use of radiocarbon from nuclear fallout as a dated marker in the otoliths of haddock *Melanogrammus aeglefinus*. *Mar. Ecol. Prog. Ser.* 1997; **150**: 49–56.
- Campana SE. Accuracy, precision and quality control in age determination, including a review of the use and abuse of age validation methods. *J. Fish. Biol.* 2001; **59**: 197–242.
- Campana SE, Zwanenburg KCT, Smith JN. $^{210}\text{Pb}/^{226}\text{Ra}$ determination of longevity in redfish. *Can. J. Fish. Aquat. Sci.* 1990; **47**: 163–165.
- Campana SE, Natanson LJ, Myklevoll S. Bomb dating and age determination of large pelagic sharks. *Can. J. Fish. Aquat. Sci.* 2002; **59**: 450–455.
- Campana SE, Jones C, McFarlane GA, Myklevoll S. Bomb dating and age validation using the spines of spiny dogfish (*Squalus acanthias*). *Env. Biol. Fishes.* 2006; **77**: 337–336.
- Caswell H. *Matrix Population Models: Construction, Analysis, and Interpretation*. 2nd ed. Sinauer Associates, Sunderland, MA. 2001; 722 pp.
- Christensen V, Guenette S, Heymans JJ, Walters CJ, Watson R, Zeller D, Pauly D. Hundred-year decline of North Atlantic predatory fishes. *Fish and Fisheries* 2003; **4**: 1–24.
- Clark M. Are deepwater fisheries sustainable?—the example of orange roughy (*Hoplostethus atlanticus*) in New Zealand. *Fish. Res.* 2001; **51**: 123–135.
- Conover DO. Nets versus nature. *Nature* 2007; **450**: 179–180.
- Conover DO, Munch SB. Sustaining fisheries yields over evolutionary time scales. *Science* 2002; **297**: 94–96.
- de Roos AM, Boukal DS, Persson L. Evolutionary regime shifts in age and size at maturation of exploited fish stocks. *Proc. Royal Soc. B* 2006; **278**: 1873–1880.
- Drazen JC, Seibel BA. Depth-related trends in metabolism of benthic and benthopelagic deep-sea fishes. *Limnol. Oceanogr.* 2007; **52**(5): 2306–2316.
- Duarte R, Azevedo M, Landa J, Pereda P. Reproduction of anglerfish (*Lophius budegassa* Spinola and *Lophius piscatorius* Linnaeus) from the Atlantic Iberian coast. *Fish. Res.* 2001; **51**: 349–361.
- Ebert T. *Plant and Animal Populations: Methods in Demography*. Academic Press, San Diego, London, Boston, New York, Sydney, Tokyo, Toronto. 1999; 312 pp.
- Edeline E, Carlson SM, Stige LC, Winfield IJ, Fletcher JM, James JB, Haugen TO, Vollestad LA, Stenseth NC. Trait changes in a harvested population are driven by a dynamic tug-of-war between natural and harvest selection. *NC. Proc. Nat. Acad. Sci.* 2007; **104**(40): 15799–15804.
- Ewing GP, Lyle JM, Murphy RJ, Kalish JM, Ziegler PE. Validation of age and growth in a long-lived temperate reef fish using otolith structure, oxytetracycline and bomb radiocarbon methods. *Mar. Freshwat. Res.* 2007; **58**: 944–955.
- Fenton GE, Short SA, Ritz DA. Age determination of orange roughy, *Hoplostethus atlanticus* (Pisces: Trachichthyidae) using $^{210}\text{Pb}/^{226}\text{Ra}$ disequilibria. *Mar. Biol.* 1991; **109**: 197–202.
- Francis RICC. The precision of otolith radiometric ageing of fish and the effect of within-sample heterogeneity. *Can. J. Fish. Aquat. Sci.* 2003; **60**: 441–447.
- Frank KT, Petrie B, Choi J, Leggett WC. Trophic cascades in a formerly cod-dominated ecosystem. *Science* 2005; **308**: 1621–1623.
- Frisk MG, Miller TJ, Dulvy, NK. Life histories and vulnerability to exploitation of elasmobranchs: inferences from elasticity, perturbation and phylogenetic analyses. *J. Northw. Atl. Fish. Sci.* 2005; **35**: 27–45.
- Gallucci VF, Saila SB, Gustafson DJ, Rothschild BJ (eds). *Stock Assessment; Quantitative Methods and Applications for Small-Scale Fisheries*. CRC/Lewis Publishers, Boca Raton, New York, London, Tokyo. 1996; 527 pp.
- Garcia VB, Lucifora LO, Myers RA. The importance of habitat and life history to extinction risk in sharks, skates, rays and chimaeras. *Proc. R. Soc. B.* 2008; **275**: 83–89.

- Gordon JDM. Deep-water fish and fisheries: introduction. Editorial. *Fish. Res.* 2001; **51**: 105–111.
- Gunderson DR, Callahan, Goiney B. Maturation and fecundity of four species of *Sebastes*. *Mar. Fish. Rev.* 1980; **42**: 74–79.
- Haedrich RL, Merrett NR, O’Dea NF. Can ecological knowledge catch up with deep-water fishing? a North Atlantic perspective. *Fish. Res.* 2001; **51**(2–3): 113–122.
- Haldorsen L, Love M. Maturity and fecundity in the rockfishes, *Sebastes* spp., a review. *Mar. Fish. Rev.* 1991; **53**(2): 25–31.
- Harvey CJ. Effects of El Niño events on energy demand and egg production of rockfish (Scorpaenidae: *Sebastes*): a bioenergetics approach. *Fish. Bull.* 2005; **103**: 71–83.
- Harvey CJ, Tolimieri N, Levin PS. Changes in body size, abundance, and energy allocation in rockfish assemblages of the northeast Pacific. *Ecol. Applic.* 2006; **16**(4): 1502–1515.
- Hauser L, Adcock GJ, Smith PJ, Bernal Ramirez JH, Carvalho GR. Loss of microsatellite diversity and low effective population size in an overexploited population of New Zealand snapper (*Pagrus auratus*). *Proc. Nat. Acad. Sci. USA.* 2002; **99**(18): 11742–11747.
- Heppell SS, Caswell H, Crowder LB. Life histories and elasticity patterns: perturbation analysis for species with minimal demographic data. *Ecology* 2000; **81**: 654–665.
- Hilborn R. Reinterpreting the state of fisheries and their management. *Ecosystems* 2007; **10**: 1362–1369.
- Hilborn R, Walters C. *Quantitative Fisheries Stock Assessment: Choice, Dynamics, and Uncertainty*. Chapman and Hall, New York. 1992.
- Hutchings JA. The cod that got away. *Nature* 2004; **428**: 899–900.
- Hutchings JA, Reynolds JD. Marine fish population collapses: consequences for recovery and extinction risk. *Bioscience* 2004; **54**: 297–309.
- Jorgensen C, Enberg K, Dunlop ES, Arlinghaus R, Boukal DS, Brander K, Ernande B, Gardmark A, Johnston F, Matsumura S, Pardoe H, Raab K, Silva A, Vainikka A, Dieckmann U, Heino M, Rijnsdorp A. Managing evolving fish stocks. *Science* 2007; **318**: 1247–1248.
- Kalish JM. Pre- and post-bomb radiocarbon in fish otoliths. *Earth Planet. Sci. Lett.* 1993; **114**: 549–554.
- Kalish JM. Use of the bomb radiocarbon chronometer to validate fish age. Final Report FRDC Project 93/109. Fisheries Research and Development Corporation, Canberra, Australia. 2001; 384 pp.
- Kalish JM, Nydal R, Nedreaas KH, Burr, GS, Eine GL. A time history of pre- and post-bomb radiocarbon in the Barents Sea derived from Arcto-Norwegian cod otoliths. *Radiocarbon* 2001; **43**(2B): 843–855.
- Kastelle CR, Kimura DK, Nevissi AE, Gunderson DR. Using Pb-210/Ra-226 disequilibria for sablefish, *Anoplopoma fimbria*, age validation. *Fishery Bull.* 1994; **92**: 292–301.
- Kastelle CR, Kimura DK, Jay SR. Using $^{210}\text{Pb}/^{226}\text{Ra}$ disequilibrium to validate conventional ages in scorpaenids (genera *Sebastes* and *Sebastolobus*). *Fish. Res.* 2000; **46**: 299–312.
- Kastelle CR, Kimura DK. Age validation of the walleye pollock (*Theragra chalcogramma*) from the Gulf of Alaska using the disequilibrium of Pb-210 and Ra-226. *ICES J. Mar. Sci.* 2006; **63**: 1520–1529.
- Kenchington E. Effets de la peche sur les especes et sur la diversite genetique—Efectos de la pesca sobre las especies y la diversidad genetica. Abstract, Reykjavik Conference on Responsible Fisheries in the Marine Ecosystem, FAO Fish. Rep. No. 658. 2002; 17–18.
- Kerr LA, Andrews AH, Frantz BR, Coale KH, Brown TA, Cailliet GM. Radiocarbon in otoliths of yelloweye rockfish (*Sebastes ruberrimus*): a reference time series for the coastal waters of southeast Alaska. *Can. J. Fish. Aquat. Sci.* 2004; **61**: 1–9.
- Kerr LA, Andrews AH, Munk K, Coale KH, Frantz BR, Cailliet GM, Brown TA. Age validation of quillback rockfish (*Sebastes maliger*) using bomb radiocarbon. *Fishery Bull.* 2005; **103**: 97–107
- Kerr LA, Andrews AH, Cailliet GM, Brown TA, Coale KH. Investigations of $\Delta^{14}\text{C}$, $\Delta^{13}\text{C}$, and $\Delta^{15}\text{N}$ in vertebrae of white shark (*Carcharodon carcharias*) from the eastern North Pacific Ocean. *Env. Biol. Fishes* 2006; **77**: 337–353.
- Kimura DK, Kastelle CR. Perspectives on the relationship between otolith growth and the conversion of isotope activity ratios to fish ages. *Can. J. Fish. Aquat. Sci.* 1995; **52**: 2296–2303.
- Kneebone J, Natanson LJ, Andrews AH, Howell WH. Using bomb radiocarbon analysis to validate age and growth estimates for the tiger shark, *Galeocerdo cuvier*, in the western North Atlantic. *Mar. Biol.* 2008 (in press).
- Landa J, Pereda P, Duarte R, Azevedo M. Growth of anglerfish (*Lophius piscatorius* and *L. budegassa*) in Atlantic Iberian waters. *Fish. Res.* 2001; **51**: 363–376.
- Lea RN, McAllister RD, VenTresca DA. Biological aspects of nearshore rockfishes of the genus *Sebastes* from central California, with notes on ecologically related sport fishes. *Fish. Bull.* 1999; **177**: 112.

- Longhurst A. Murphy's law revisited: longevity as a factor in recruitment to fish populations. *Fish. Res.* 2002; **56**: 125–131.
- Longhurst A. The sustainability myth. *Viewpoint In Fish. Res.* 2006; **81**: 107–1121.
- Love MS, Morris P, McCrae M, Collins R. Life history aspects of 19 rockfish species (Scorpaenidae: *Sebastes*) from the southern California Bight. NOAA Tech. Report NMFS. No. 87. 1990; 38 pp.
- Love MS, Yoklavich MM, Thorsteinson L. *The Rockfishes of the Northeast Pacific*. University of California Press, Berkeley, Los Angeles, London. 2002; 404 pp.
- MacCall AD. Status of bocaccio off California in 2007. In: Status of the Pacific coast groundfish fishery through 2003, stock assessment and fishery evaluation. Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 200, Portland, OR 97220-1384. 2007.
- Marriott PM, Manning MJ, Horn PL. Age estimation and maturity of the ridge-scaled macrourid (*Macrourus whitsoni*) from the Ross Sea. *CCAMLR Sci.* 2006; **13**: 1–13.
- Mason J. Declining rockfish lengths in the Monterey Bay, California, 1959–94. *Mar. Fish. Rev.* 1998; **60**(3): 15–28.
- Methot RD. Synthesis model: an adaptable framework for analysis of diverse stock assessment data. *INPFC Bull.* 1990; **50**: 259–277.
- Methot R, Piner K. Status of the canary rockfish resource off California, Oregon and Washington in 2001. STAR Panel Review, Pacific Fishery Management Council. 2002; 1–8.
- Millar RB, Methot RD. Age-structured meta-analysis of U.S. West Coast rockfish (Scorpaenidae) populations and hierarchical modeling of trawl survey catchabilities. *Can. J. Fish. Aquat. Sci.* 2002; **59**: 383–392.
- Milton DA, Short SA, O'Neill MF, Blaber SJM. Ageing of three species of tropical snapper (Lutjanidae) from the Gulf of Carpentaria, Australia, using radiometry and otolith ring counts. *Fish. Bull.* 1995; **93**: 103–115.
- Mollet HF, Cailliet GM. Comparative population demography of elasmobranchs using life history tables, Leslie matrices, and stage-based matrix models. *Mar. Freshwat. Res.* 2002; **53**(2): 503–516.
- Mollet HF, Cailliet GM. Reply to comments by Miller *et al.* on Mollet and Cailliet (2002): confronting models with data. *Mar. Freshwat. Res.* 2003; **54**: 739–744.
- Munk K. Maximum ages of groundfishes in waters off Alaska and British Columbia and considerations of age determination. *Alaska Fish. Res. Bull.* 2001; **8**(1): 12–21.
- Myers RA, Worm B. Rapid world-wide depletion of predatory fish communities. *Nature* (Lond.) 2003; **423**: 280–283.
- Myers RA, Worm B. Extinction, survival or recovery of large predatory fishes. *Phil. Trans. R. Soc. B* 2005; **360**: 13–20.
- Myers RA, Myers JK, Shepherd TD, Powers SP, Peterson CH. Cascading effects of the loss of apex predatory sharks from a coastal ocean. 2003. *Science* 2003; **315**: 1846–1850.
- National Research Council. *Marine Protected Areas: Tools for Sustaining Ocean Ecosystems*. National Academy Press, Washington DC. 2001.
- Novikov GG, Shibanov VN, Stroganov AN, Afanasjev KI, Melnikov SP. Analysis of population and genetic characteristics of redfish of the Irminger Sea. Poster. ICES CM 2002; 10.
- O'Farrell MR, Botsford LW. Estimation of change in lifetime egg production from length frequency data. *Can. J. Fish. Aquat. Sci.* 2005; **62**: 1626–1639.
- O'Farrell MR, Botsford LW. The fisheries management implications of maternal-age-dependent larval survival. *Can. J. Fish. Aquat. Sci.* 2006a; **63**: 2249–2258.
- O'Farrell MR, Botsford LW. Estimating the status of nearshore rockfish (*Sebastes* spp.) populations with length frequency data. *Ecol. Applic.* 2006b; **16**: 977–986.
- Olsen EM, Heino M, Lilly GR, Morgan MJ, Brattey J, Ernande B, Dieckmann U. Maturation trends indicative of rapid evolution preceded the collapse of northern cod. *Nature* 2004; **428**: 932–935.
- Pankhurst, NW, Conroy AM. Size-fecundity relationships in the orange roughy, *Hoplostethus atlanticus*. *New Zeal. J. Mar. Freshwat. Res.* 1987; **21**: 295–300.
- Parker SJ, Berkeley SA, Golden JT, Gunderson DR, Heifetz J, Hixon MA, Larson R, Leaman BM, Love MS, Musick JA, O'Connell VM, Ralston S, Weeks HF, Yoklavich MM. Management of Pacific rockfish. *AFS Policy Statement Fisheries* 2000; **25**(3): 22–29.
- Pauly D. Anecdotes and the shifting baseline syndrome of fisheries. *Trends Ecol. Evol.* 1995; **10**: 430.
- Pauly D, Christensen V. Primary production required to sustain global fisheries. *Nature* 1995; **374**: 255–257.
- Pauly D, Christensen V, Dalsgaard J, Froese R, Torres F Jr. Fishing down marine food webs. *Science* 1998; **279**: 860–863.

- Pauly D, Christensen V, Guenette S, Pitcher TJ, Sumaila UR, Walters CJ, Watson R, Zeller D. Towards sustainability in world fisheries. *Nature* 2002; **418**: 689–695.
- Pauly D, Alder J, Bennett E, Christensen V, Tyedmers P, Watson R. The future for fisheries. *Science* 2003; **302**: 1359–1361.
- Phillips JB. Life history studies on ten species of rockfish (genus *Sebastes*). *Fish. Bull.* 1964; **126**: 1–70.
- Piner, KR, Hamel OS, Henkel JL, Wallace JR, Hutchinson CE. Age validation of canary rockfish (*Sebastes pinniger*) from off the Oregon coast (USA) using the bomb radiocarbon method. *Can. J. Fish. Aquat. Sci.* 2005; **62**: 1060–1066.
- Piner KR, Wischniowski SG. Pacific halibut (*Hippoglossus stenolepis*) chronology of bomb radiocarbon in otoliths from 1944–1981, and a validation of ageing methods. *J. Fish Biol.* 2004; **64**: 1060–1071.
- Ralston S, Bence JR, Eldridge MB, Lenarz WH. An approach to estimating rockfish biomass based on larval production, with application to *Sebastes jordani*. *Fish. Bull.* 2003; **101**: 129–146.
- Roberts CM. Deep impact: the rising toll of fishing in the deep sea. *Trends Ecol. Evol.* 2002; **17**: 242–245.
- Rogers AD, Morley S, Fitzcharles E, Jarvis K, Belchier M. Genetic structure of Patagonian toothfish (*Dissostichus eleginoides*) populations on the Patagonian Shelf and Atlantic and western Indian Ocean sections of the Southern Ocean. *Mar. Bio.* 2006; **149**: 915–924.
- Safina C, Rosenberg AA, Myers RA, Quinn TJ II, Collie JS. U.S. Ocean fish recovery: staying the course. *Science* 2005; **309**: 707–708.
- Scheffer M, Carpenter S, deYoung B. Cascading effects of overfishing marine systems. *Trends Ecol. Evol.* 2005; **20**(11): 579–581.
- Sibert J, Hampton J, Kleiber P, Maunder M. Biomass, Size, and trophic status of top predators in the Pacific Ocean. *Science* 2006; **314**: 1773–1776.
- Sigurdsson T, Thorsteinsson V, Gustafsson L. *In situ* tagging of deep-sea redfish: application of an underwater, fish-tagging system. *ICES J. Mar. Sci.* 2006; **63**: 523–531.
- Smith DC, Fenton GE, Robertson SG, Short SA. Age determination and growth of orange roughy (*Hoplostethus atlanticus*): a comparison of annulus counts with radiometric ageing. *Can. J. Fish. Aquat. Sci.* 1995; **52**: 391–401.
- Smith JN, Nelson R, Campana SE. The use of Pb-210/Ra-226 and Th-228/Ra-228 disequilibria in the ageing of otoliths of marine fish. In: Kershaw PJ, Woodhead DS (eds). *Radionuclides in the Study of Marine Processes*. Elsevier Applied Science, NewYork. 1991; 350–359.
- Starr RM, Heine JN, Felton JM, Cailliet GM. Movements of bocaccio (*Sebastes paucispinis*) and greenspotted (*S. chlorostictus*) rockfishes in a Monterey submarine canyon: implications for the design of marine reserves. *Fish. Bull.* 2002; **100**(2): 324–337.
- Stevens MM, Andrews AH, Cailliet GM, Coale KH, Lundstrom CC. Radiometric validation of age, growth, and longevity for the blackgill rockfish (*Sebastes melanostomus*). *Fishery Bull.* 2004; **102**: 711–722.
- Stewart BD, Fenton GE, Smith DC, Short SA. Validation of otolith-increment age estimates for a deepwater fish species, the warty oreo *Alloctytus verrucosus*, by radiometric analysis. *Mar. Biol.* 1995; **123**: 29–38.
- Stewart IJ, Piner KR. Simulation of the estimation of ageing bias inside an integrated assessment of canary rockfish using age estimates from a bomb radiocarbon study. *Mar. Freshwat. Res.* 2007; **58**: 905–913.
- Stockley B, Menezes G, Pinho MR, Rogers AD. Genetic population structure in the black-spot sea bream (*Pagellus bogaraveo* Brunnich, 1768) from the NE Atlantic. *Mar. Biol.* 2005; **146**: 793–804.
- Stransky C, Kanisch G, Kruger A, Purkl S. Radiometric age validation of golden redfish (*Sebastes marinus*) and deep-sea redfish (*S. mentella*) in the Northeast Atlantic. *Fish. Res.* 2005; **74**: 186–197.
- VenTresca DA, Parrish RH, Houk JL, Gingras ML, Short SD, Crane NL. El Niño effects on the somatic and reproductive condition of blue rockfish, *Sebastes mystinus*. *Cal. Coop. Oceanic Fish. Invest. Rep.* 1995; **36**: 167–174.
- Walsh MR, Munch S, Chiba S, Conover DO. Maladaptive changes in multiple traits caused by fishing: impediments to population recovery. *Ecology Lett.* 2006; **9**: 142–148.
- Walters C. Folly and fantasy in the analysis of spatial catch rate data. *Can. J. Fish. Aquat. Sci.* 2003; **60**: 1433–1436.
- Watters DL, Kline DE, Coale KH, Cailliet GM. Radiometric age confirmation and growth of bank rockfish, *Sebastes rufus*, from California. *Fish. Res.* 2006; **81**: 251–257.
- Welden BA, Cailliet GM, Flegal AR. Comparison of radiometric vertebral band age estimates in four California elasmobranchs. In: Summerfelt RC, Hall GE (eds). *The Age and Growth of Fish*. The Iowa State University Press, Ames. 1987; 301–315.

- Worm B, Sandow M, Oschlies A, Lotze HK, Myers RA. Global patterns of predator diversity in the open oceans. *Science* 2005; **309**: 1365–1369.
- Ziegler PE, Lyle JM, Haddon M, Ewing GP. Rapid changes in life-history characteristics of a long-lived temperate reef fish. *Mar. Freshwat. Res.* 2007; **58**: 1096–1107.